

The status of taxa of subgenus *Aedes* (Diptera, Culicidae, *Aedes*): *esoensis* Yamada, *rossicus* Dolbeskin, Gorickaja & Mitrofanova, *cinereus* Meigen, *geminus* Peus. II. Illustrations

Rimma M. Gornostaeva

26 Bakinskih komissrov str., 3-1-447, 117571 Moscow, Russia. Email: gornostaeva@newmail.ru

Introduction / Abstract

Illustrations are given of the species *cinereus*, *esoensis*, *rossicus* and *geminus* of subgenus *Aedes* to accompany the morphological descriptions given in the first part of this article, which appeared in *European Mosquito Bulletin* 16, 22-26.

Legends to figures

Figure 1. Measurement of male genitalia.

Figure 2. Genitalia of the male *Aedes rossicus* from Perm Region (male from type-series); ventral (1) and dorsal (2) aspects.

Figure 3. *Aedes rossicus*. Ventral (1) and dorsal (2) aspects of male from Tatar Republic. Ventral aspect of male from Krasnoyarsk Territory (3).

Figure 4. *Aedes rossicus*. Claspettes of males from Tatar Republic (1), Khakassia (2), Perm Region (3), Buriatia (4), Tuva (5). Apical parts of lateral branches of gonostylus of males from Tatar Republic (6), Krasnoyarsk Territory (8), Khakassia (9), Buriatia (7, 10, 11). The lobes of IX tergum of the male from Buriatia (14). Medial branch of gonostylus of male from Tatar Republic (18). Claspettes (12, 13, 15, 16, 17, 19-21) of males from Buriatia (Taximo).

Figure 5. Basal mesal lobes of *Aedes cinereus* from Moscow Region (1), *Aedes rossicus* from Krasnoyarsk Territory (2), *Aedes geminus* from Moscow Region (3) and *Aedes rossicus* from Maritime Territory (= Primorski Krai) (4).

Figure 6. *Aedes esoensis*. Ventral (1) and dorsal (2) aspects. Claspette (7), medial branch of gonostylus (8), apical part of the lateral branch of gonostylus (9), lobe of tergum IX (10) of male from Yuzhno-Sakhalinsk. Apical parts of the lateral branches of gonostylus in different positions of males from Yuzhno-Sakhalinsk (3, 5); Maritime Territory (4, 6). Claspettes of males from Yuzhno-Sakhalinsk (11, 12, 15) and Maritime Territory (13, 14).

Figure 7. *Aedes cinereus*. Ventral aspect of newly emerged male (1), dorsal aspect of newly emerged male (2), ventral aspect of male prepared for copulation (3). The apical parts of the lateral branch of the gonostylus of males from Moscow Region (4-10), Belarus (11), Ural (12). Claspettes of males from Moscow Region (13-15, 18), Ukraine (19), Belarus (20), the lobe of tergum IX (16) and medial branch of gonostylus (17) of male from Moscow Region.

Figure 8. *Aedes geminus* from Moscow Region. Ventral aspect (1), dorsal aspect (2), ventral aspect of newly emerged male (3), claspettes (4-6) and aspects of gonostylus (7-9).

Figure 9. *Aedes geminus*. Ventral (1) and dorsal (2) aspects of male from Chita Region. Apical parts of the lateral branch of gonostylus (3), claspette (4), medial branch of gonostylus (5, 6). Apical parts of lateral branch of gonostylus of males from Moscow Region (7, 8), Khakassia (9, 11), Buriatia (10), Tuva (12, 13), Maritime Territory (14, 15). Claspettes of males from Belarus (16), Moscow Region (17), Khakassia (18, 19), Tuva (20, 21), Buriatia (22, 23). Lobes of tergum IX of males from Moscow Region (24) and Tuva (25).




Figure 1





Figure 2




Figure 3




Figure 4



Figure 5







Figure 6









Figure 7




Figure 8




Figure 9